

| Patient Information | | | | Prescriber Information | |
|--|---------|------------------|--|--|-----------|
| Patient Name: | | DOB: | | Prescriber's Name: | |
| Address: | | | | NPI#: | |
| City: | State: | Zip: | | DEA#: | License#: |
| Phone: | | Alternate Phone: | | SSN: | |
| Height: | Weight: | Allergies: | | Phone: | Fax: |
| Emergency Contact: | | Phone: | | Contact Person: | |
| Is the patient pregnant, planning a pregnancy or nursing: Yes No | | | | Does the patient need interpreter services: Yes No | |

| Documentation (Required) | | |
|---------------------------------|---------------------|------------------------|
| Insurance Card (front and back) | Current Medications | History/Progress Notes |

Specific Labs Needed (note: send ALL labs, must include specific labs listed here)

Other: _____

| Clinical Information | |
|-------------------------|--|
| Primary Diagnosis _____ | <i>**Please include Dx Code # and description Prior Failed</i> |
| Meds: _____ | |

| Prescription Information | | | |
|---------------------------------|--------------|------------------------------|-----------|
| Patient Medical History | ICD-10 Code: | Prior Failed Medications: | Duration: |
| Date of Osteoporosis diagnosis: | | Fosamax (alendronate) | |
| DEXA T-score (worst sites): | | Actonel (risdrionate) | |
| Previous Fracture(s) Yes No | | Miacalcin Nasal Spray | |
| Site of Fracture(s): | | Boniva | |
| Others: | | Reclast | |

| Medication | Dose | Directions for Use | Qty | Refills |
|----------------------------|------------------|---|--|---------|
| FORTEO | 600mcg/2.4mL Pen | Inject 1 dose (20mcg) subcutaneously once daily. Discard device 28 days after first use. | 1 pen (4 -week supply) 3 pens(12 week supply) | |
| BD MINI PEN NEEDLES | 31Gx3/16" | Use with Forteo pen once daily as directed | #90 pen needles #30 pen needles | |
| PROLIA | 60mg/1mL vial | Inject the contents of 1 syringe (60mg) subcutaneously every 6 months | 1 prefilled Syringe | |
| RECLAST | 5mg/100mL vial | Infuse 5mg intravenously over no less than 15 minutes once annually. | One: 5mg/100mL vial | |
| BONIVA | 3mg/3mL PFS | Inject the contents of 1 syringe (3mg) intravenously every 3 months. To be administrated by a healthcare professional | One: 3mg/3mLPFS | |

Patient has received pen and injection training

Physician's office to provide injection training

By signing this form I authorize AleraCare and its representatives to act as my agent in order to initiate and execute the insurance prior authorization process and, in doing so release clinical information via phone to the appropriate third party payer.

 Prescriber's Signature

 Date