

**Patient Information**

Patient Name:		DOB:				
Patient Home Phone:	Patient Cell Phone:	Patient Email:				
Emergency/Alternate Contact Name:		Emergency/Alternate Contact Phone:				
NKDA	Allergies:	Weight lbs/kg:	Height:			
Patient Status:	New to Therapy	Continuing Therapy	Therapy Change	Last infusion date (if applicable): _____		
Is the patient pregnant, planning a pregnancy or nursing:		Yes	No	Does the patient need interpreter services:	Yes	No

**Provider Information**

Referral Coordinator Name:		Referral Coordinator Email:		
Ordering Provider:		Provider NPI:		
Referring Practice Name:		Phone:	Fax:	
Practice Address:		City:	State:	Zip:

**ICD-10 CODE**

G35 Multiple Sclerosis                      Other: \_\_\_\_\_

**Medication Order**

<b>Briumvi</b> (ublituximab-xiiv)	<b>Initiation Therapy:</b>	<b>Maintenance Therapy:</b>
	150mg IV for first infusion    Qty: 1 vial	450mg IV every 24 weeks    Qty:3 vials
	450mg IV 2 weeks after the first infusion    Qty:3 vials	
	450mg IV 24 weeks after the first infusion and then 24 weeks thereafter    Qty:3 vials	

**Documentation Required (Note: Send all labs, must include specific labs listed here)**

Labs: Hepatitis B; Quantitative Serum immunoglobulins	Insurance Card (front and back)
Current Medications	History/Progress Notes

**Pre-Medication Order**

<b>methylprednisolone</b> (Required; give 30min before infusion)	100mg IV	<b>loratadine</b> (Claritin)	10mg PO
<b>acetaminophen</b> (Tylenol)	500mg    650mg    1000mg PO	<b>diphenhydramine</b> (Benadryl)	25mg IV    50mg IV
<b>cetirizine</b> (Zyrtec)	10mg PO		
Other: _____			
Dose: _____	Route: _____	Frequency: _____	

**Line/Port Use Orders**

RN to start/discontinue peripheral IV or access/de-access port

RN to administer 3-10 mL normal saline with heparin 100 units/mL 1-5 mL per SASH method

Do NOT administer heparin to this patient

Additional orders: \_\_\_\_\_

**Adverse Reaction and Anaphylaxis Orders**

Standard anaphylaxis kit (epinephrine 0.3 mg IM PRN for severe reactions x 1, may repeat once within 5-15 minutes for a maximum of 2 doses; Benadryl 50 mg PO PRN for mild-moderate reactions x 1; Benadryl 50 mg IV PRN for mild-moderate reactions x 1)

Additional orders: \_\_\_\_\_

**Special Instructions (Prior therapy, treatment dates, and reasons for d/c)****Provider Name**

Order is valid for 1 year from date of signature and refills will be provided to cover the duration of treatment unless otherwise indicated.

**Provider Signature****Date**

Check here if this is a stat order