

Patient Information

Patient Name:		DOB:	
Patient Home Phone:	Patient Cell Phone:	Patient Email:	
Emergency/Alternate Contact Name:		Emergency/Alternate Contact Phone:	
NKDA	Allergies:	Weight lbs/kg:	Height:
Patient Status:	New to Therapy	Continuing Therapy	Therapy Change
Is the patient pregnant, planning a pregnancy or nursing:		Yes	No
Does the patient need interpreter services:		Yes	No

Provider Information

Referral Coordinator Name:	Referral Coordinator Email:		
Ordering Provider:	Provider NPI:		
Referring Practice Name:	Phone:	Fax:	
Practice Address:	City:	State:	Zip:

ICD-10 CODE

K51.80 Ulcerative colitis	K51.90 Ulcerative colitis, unspecified, without complications
K50.90 Crohn's disease, unspecified, without complications	K50.00 Crohn's disease of small intestine without complications
K50.10 Crohn's disease of large intestine without complications	Other: _____

Medication Order

Entyvio (vedolizumab)	Dose: 300mg IV over 30 min	Frequency: Week 0, 2, 6 and then every 8 weeks thereafter	Qty: 3 doses
		Every 8 weeks	Qty: 1 dose

Documentation Required (Note: Send all labs, must include specific labs listed here)

Labs (TB test)	Insurance Card (front and back)	Current Medications	History/Progress Notes
----------------	---------------------------------	---------------------	------------------------

Pre-Medication Order

acetaminophen (Tylenol)	500mg	650mg	1000mg PO	diphenhydramine (Benadryl)	25mg	50mg /	PO	IV
cetirizine (Zyrtec)	10mg PO			methylprednisolone (Solu-Medrol)	40mg IV		125mg IV	
loratadine (Claritin)	10mg PO			hydrocortisone (Solu-Cortef)	100mg IV			
Other: _____								
Dose: _____			Route: _____			Frequency: _____		

Line/Port Use Orders

RN to start/discontinue peripheral IV or access/de-access port

RN to administer 3-10 mL normal saline prior to medication and 30 mL normal saline after medication with heparin 100 units/mL 1-5 mL per SASH method

Do NOT administer heparin to this patient

Additional orders: _____

Adverse Reaction and Anaphylaxis Orders

Standard anaphylaxis kit (epinephrine 0.3 mg IM PRN for severe reactions x 1, may repeat once within 5-15 minutes for a maximum of 2 doses; Benadryl 50 mg PO PRN for mild-moderate reactions x 1; Benadryl 50 mg IV PRN for mild-moderate reactions x 1)

Additional orders: _____

Special Instructions (Prior therapy, treatment dates, and reasons for d/c)
Provider Name

Order is valid for 1 year from date of signature and refills will be provided to cover the duration of treatment unless otherwise indicated.

Provider Signature
Date

Check here if this is a stat order