

**Patient Information**

Patient Name:		DOB:	
Patient Home Phone:	Patient Cell Phone:	Patient Email:	
Emergency/Alternate Contact Name:		Emergency/Alternate Contact Phone:	
NKDA	Allergies:	Weight lbs/kg:	Height:
Patient Status:	New to Therapy	Continuing Therapy	Therapy Change
Is the patient pregnant, planning a pregnancy or nursing:		Last infusion date (if applicable): _____	
Yes	No	Does the patient need interpreter services:	Yes No

**Provider Information**

Referral Coordinator Name:	Referral Coordinator Email:		
Ordering Provider:	Provider NPI:		
Referring Practice Name:	Phone:	Fax:	
Practice Address:	City:	State:	Zip:

**ICD-10 CODE**

D69.3 Idiopathic thrombocytopenia purpura (ITP)	G61.81 Chronic inflammatory demyelinating polyneuropathy (CIDP)
D80.9 Primary humoral immunodeficiency (PI)	D83.9 Common variable immunodeficiency/agammaglobulinemia
D82.0 Wiskott-Aldrich syndrome	G61.82 Multifocal motor neuropathy
	M33.13 Dermatomyositis without myopathy
Other: _____	

**Documentation Required (Note: Send all labs, must include specific labs listed here)**

Labs (IgG)	Insurance Card (front and back)	Current Medications	History/Progress Notes
------------	---------------------------------	---------------------	------------------------

**Medication Order**

Gammagard 10%	Privigen 10%
---------------	--------------

**Dose and frequency:**

\_\_\_\_\_ gm/kg or \_\_\_\_\_ grams IV divided over \_\_\_\_\_ days every \_\_\_\_\_ weeks

\_\_\_\_\_ grams IV every \_\_\_\_\_ weeks

\_\_\_\_\_ mg/kg IV every \_\_\_\_\_ weeks

Other: \_\_\_\_\_

**Note:** Pharmacist will calculate infusion rates unless otherwise specified: \_\_\_\_\_

**Pre-Medication Order**

<b>acetaminophen</b> (Tylenol)	500mg	650mg	1000mg PO	<b>diphenhydramine</b> (Benadryl)	25mg	50mg /	PO	IV
<b>cetirizine</b> (Zyrtec)	10mg PO			<b>methylprednisolone</b> (Solu-Medrol)	40mg IV	125mg IV		
<b>loratadine</b> (Claritin)	10mg PO			<b>hydrocortisone</b> (Solu-Cortef)	100mg IV			
Other: _____								
Dose: _____			Route: _____			Frequency: _____		

**Line/Port Use Orders**

RN to start/discontinue peripheral IV or access/de-access port

RN to administer 3-10 mL normal saline with heparin 100 units/mL 1-5 mL per SASH method

Do NOT administer heparin to this patient

Additional orders: \_\_\_\_\_

**Adverse Reaction and Anaphylaxis Orders**

Standard anaphylaxis kit (epinephrine 0.3 mg IM PRN for severe reactions x 1, may repeat once within 5-15 minutes for a maximum of 2 doses;  
Benadryl 50 mg PO PRN for mild-moderate reactions x 1; Benadryl 50 mg IV PRN for mild-moderate reactions x 1)

Additional orders: \_\_\_\_\_

**Special Instructions (Prior therapy, treatment dates, and reasons for d/c)**

\_\_\_\_\_  
**Provider Name**

Order is valid for 1 year from date of signature and refills will be provided to cover the duration of treatment unless otherwise indicated.

\_\_\_\_\_  
**Provider Signature**

\_\_\_\_\_  
**Date**

Check here if this is a stat order