aleracare
advanced home infusion

FAX 818.390.9697

Orencia (abatacept)

Provider Order Form rev. 10/4/24

Patient Information

Patient Name: DOB:

Patient Home Phone: Patient Cell Phone: Patient Email:

Emergency/Alternate Contact Name: Emergency/Alternate Contact Phone:

◯ NKDA ◯ Allergies: Weight lbs/kg: Height:

Patient Status: ◯ New to Therapy ◯ Continuing Therapy ◯ Therapy Change Last infusion date (if applicable): _____

Is the patient pregnant, planning a pregnancy or nursing: ◯ Yes ◯ No Does the patient need interpreter services: ◯ Yes ◯ No

Provider Information

Referral Coordinator Name: Referral Coordinator Email:

Ordering Provider: Provider NPI:

Referring Practice Name: Phone: Fax:

Practice Address: City: State: Zip:

ICD-10 CODE

◯ M06.9 Rheumatoid arthritis ◯ L40.50 Psoriatic Arthritis ◯ M08.40 Polyarticular juvenile idiopathic arthritis

◯ D89.813 Graft versus host disease, unspecified (aGVHD) ◯ Other: _____

Medication Order

Orencia (abatacept) **Dose:** ◯ 500mg IV ◯ 750mg IV ◯ 1,000mg IV ◯ 10mg/kg (____mg) (maximum of 1,000mg)

IV INFUSION ◯ Other: _____

Frequency: ◯ Initiation therapy: Administer at 0, 2 and 4 weeks then every 4 weeks thereafter Qty: 3 doses

◯ Maintenance: Every 4 weeks Qty: 1 dose

◯ aGVHD: Infuse over 60 minutes on the day before transplantation, followed by a dose on Day 5, 14 and 28 after transplant Qty: 4 doses

Documentation Required (Note: Send all labs, must include specific labs listed here)

◯ Labs (TB and Hep B) ◯ Insurance Card (front and back) ◯ Current Medications ◯ History/Progress Notes

Pre-Medication Order

acetaminophen (Tylenol) ◯ 500mg ◯ 650mg ◯ 1000mg PO

cetirizine (Zyrtec) ◯ 10mg PO

loratadine (Claritin) ◯ 10mg PO

diphenhydramine (Benadryl) ◯ 25mg ◯ 50mg / ◯ PO ◯ IV

methylprednisolone (Solu-Medrol) ◯ 40mg IV ◯ 125mg IV

hydrocortisone (Solu-Cortef) ◯ 100mg IV

Other: _____

Dose:_____ Route:_____ Frequency: _____

Line/Port Use Orders

RN to start/discontinue peripheral IV or access/de-access port

◯ RN to administer 3-10 mL normal saline with heparin 100 units/mL 1-5 mL per SASH method

◯ Do NOT administer heparin to this patient

◯ Additional orders: _____

Adverse Reaction and Anaphylaxis Orders

◯ Standard anaphylaxis kit (epinephrine 0.3 mg IM PRN for severe reactions x 1, may repeat once within 5-15 minutes for a maximum of 2 doses; Benadryl 50 mg PO PRN for mild-moderate reactions x1; Benadryl 50 mg IV PRN for mild-moderate reactions x1)

◯ Additional orders: _____

Special Instructions (Prior therapy, treatment dates, and reasons for d/c)

Provider Name Provider Signature Date

Order is valid for 1 year from date of signature and refills will be provided to cover the duration of treatment unless otherwise indicated.

Check here if this is a stat order ◯