

**Patient Information**

Patient Name:		DOB:	
Patient Home Phone:	Patient Cell Phone:	Patient Email:	
Emergency/Alternate Contact Name:		Emergency/Alternate Contact Phone:	
NKDA	Allergies:	Weight lbs/kg:	Height:
Patient Status:	New to Therapy	Continuing Therapy	Therapy Change
Last infusion date (if applicable):		_____	
Is the patient pregnant, planning a pregnancy or nursing:		Yes	No
Does the patient need interpreter services:		Yes	No

**Provider Information**

Referral Coordinator Name:		Referral Coordinator Email:	
Ordering Provider:		Provider NPI:	
Referring Practice Name:		Phone:	Fax:
Practice Address:		City:	State: Zip:

**ICD-10 CODE**

D69.3 Idiopathic thrombocytopenia purpura (ITP)      G61.81 Chronic inflammatory demyelinating polyneuropathy (CIDP)  
 D80.9 Primary humoral immunodeficiency (PI)      D83.9 Common variable immunodeficiency/agammaglobulinemia  
 D82.0 Wiskott-Aldrich syndrome      G61.82 Multifocal motor neuropathy      M33.13 Dermatomyositis without myopathy  
 Other: \_\_\_\_\_

**Documentation Required (Note: Send all labs, must include specific labs listed here)**

Labs (IgG)	Insurance Card (front and back)	Current Medications	History/Progress Notes
------------	---------------------------------	---------------------	------------------------

**Medication Order**

Gammagard 10%      Privigen 10%

**Dose and frequency:**

\_\_\_\_\_ gm/kg or \_\_\_\_\_ grams IV divided over \_\_\_\_\_ days every \_\_\_\_\_ weeks  
 \_\_\_\_\_ grams IV every \_\_\_\_\_ weeks  
 \_\_\_\_\_ mg/kg IV every \_\_\_\_\_ weeks  
 Other: \_\_\_\_\_

**Note:** Pharmacist will calculate infusion rates unless otherwise specified: \_\_\_\_\_

**Pre-Medication Order (given 30 min before each infusion)**

<b>acetaminophen</b> (Tylenol)	500mg	650mg	1000mg PO	<b>diphenhydrAMINE</b> (Benadryl)	25mg	50mg /	PO	IV
<b>cetirizine</b> (ZyrTEC)	10mg PO			<b>methylPREDNISolone</b> (SOLU-Medrol)	40mg IV	125mg IV		
<b>loratadine</b> (Claritin)	10mg PO			<b>hydrocortisone</b> (Solu-CORTEF)	100mg IV			

Other: \_\_\_\_\_

Dose: \_\_\_\_\_      Route: \_\_\_\_\_      Frequency: \_\_\_\_\_

**Line/Port Use Orders**

RN to start/discontinue peripheral IV or access/de-access port  
 RN to administer 3-10 mL normal saline with heparin 100 units/mL 1-5 mL per SASH method  
 Do NOT administer heparin to this patient  
 Additional orders: \_\_\_\_\_

**Adverse Reaction and Anaphylaxis Orders**

Standard anaphylaxis kit (EPINEPHrine 0.3 mg IM PRN for severe reactions x 1, may repeat once within 5-15 minutes for a maximum of 2 doses;  
Benadryl 50 mg PO PRN for mild-moderate reactions x 1; Benadryl 50 mg IV PRN for mild-moderate reactions x 1)

Additional orders: \_\_\_\_\_

**Special Instructions (Prior therapy, treatment dates, and reasons for d/c)**

\_\_\_\_\_  
**Provider Name**

Order is valid for 1 year from date of signature and refills will be provided to cover the duration of treatment unless otherwise indicated.

\_\_\_\_\_  
**Provider Signature**

\_\_\_\_\_  
**Date**

Check here if this is a stat order