

**Patient Information**

Patient Name:		DOB:	
Patient Home Phone:	Patient Cell Phone:	Patient Email:	
Emergency/Alternate Contact Name:		Emergency/Alternate Contact Phone:	
NKDA	Allergies:	Weight lbs/kg:	Height:
Patient Status:	New to Therapy	Continuing Therapy	Therapy Change
Is the patient pregnant, planning a pregnancy or nursing:		Last infusion date (if applicable): _____	
Yes	No	Does the patient need interpreter services:	Yes No

**Provider Information**

Referral Coordinator Name:	Referral Coordinator Email:		
Ordering Provider:	Provider NPI:		
Referring Practice Name:	Phone:	Fax:	
Practice Address:	City:	State:	Zip:

**ICD-10 CODE**

K51.80 Ulcerative colitis	K51.90 Ulcerative colitis, unspecified, without complications
K50.90 Crohn's disease, unspecified, without complications	K50.00 Crohn's disease of small intestine without complications
K50.10 Crohn's disease of large intestine without complications	Other: _____

**Medication Order**

<b>Entyvio (vedolizumab)</b>	<b>Dose:</b> 300mg IV over 30 min	<b>Frequency:</b> Week 0, 2, 6 and then every 8 weeks thereafter	Qty: 3 doses
		Every 8 weeks	Qty: 1 dose

**Documentation Required (Note: Send all labs, must include specific labs listed here)**

Labs (TB test)	Insurance Card (front and back)	Current Medications	History/Progress Notes
----------------	---------------------------------	---------------------	------------------------

**Pre-Medication Order (given 30 min before each infusion)**

<b>acetaminophen</b> (Tylenol)	500mg	650mg	1000mg PO	<b>diphenhydrAMINE</b> (Benadryl)	25mg	50mg /	PO	IV
<b>cetirizine</b> (ZyrTEC)	10mg PO			<b>methylPREDNISolone</b> (SOLU-Medrol)	40mg IV		125mg IV	
<b>loratadine</b> (Claritin)	10mg PO			<b>hydrocortisone</b> (Solu-CORTEF)	100mg IV			

Other: \_\_\_\_\_

Dose: \_\_\_\_\_ Route: \_\_\_\_\_ Frequency: \_\_\_\_\_

**Line/Port Use Orders**

RN to start/discontinue peripheral IV or access/de-access port

RN to administer 3-10 mL normal saline prior to medication and 30 mL normal saline after medication with heparin 100 units/mL 1-5 mL per SASH method

Do NOT administer heparin to this patient

Additional orders: \_\_\_\_\_

**Adverse Reaction and Anaphylaxis Orders**

Standard anaphylaxis kit (EPINEPHrine 0.3 mg IM PRN for severe reactions x 1, may repeat once within 5-15 minutes for a maximum of 2 doses; Benadryl 50 mg PO PRN for mild-moderate reactions x 1; Benadryl 50 mg IV PRN for mild-moderate reactions x 1)

Additional orders: \_\_\_\_\_

**Special Instructions (Prior therapy, treatment dates, and reasons for d/c)**
**Provider Name**

Order is valid for 1 year from date of signature and refills will be provided to cover the duration of treatment unless otherwise indicated.

**Provider Signature**
**Date**

Check here if this is a stat order