

Patient Information

Patient Name:		DOB:				
Patient Home Phone:	Patient Cell Phone:	Patient Email:				
Emergency/Alternate Contact Name:		Emergency/Alternate Contact Phone:				
NKDA	Allergies:	Weight lbs/kg:	Height:			
Patient Status:	New to Therapy	Continuing Therapy	Therapy Change	Last infusion date (if applicable): _____		
Is the patient pregnant, planning a pregnancy or nursing:		Yes	No	Does the patient need interpreter services:	Yes	No

Provider Information

Referral Coordinator Name:	Referral Coordinator Email:		
Ordering Provider:	Provider NPI:		
Referring Practice Name:	Phone:	Fax:	
Practice Address:	City:	State:	Zip:

ICD-10 CODE

M06.9 Rheumatoid arthritis L40.50 Psoriatic Arthritis M08.40 Polyarticular juvenile idiopathic arthritis
D89.813 Graft versus host disease, unspecified (aGVHD) Other: _____

Medication Order

Orencia (abatacept) **Dose:** 500mg IV 750mg IV 1,000mg IV 10mg/kg (____mg) (maximum of 1,000mg)
IV INFUSION Other: _____

Frequency: Initiation therapy: Administer at 0, 2 and 4 weeks then every 4 weeks thereafter Qty: 3 doses
 Maintenance: Every 4 weeks Qty: 1 dose

aGVHD: Infuse over 60 minutes on the day before transplantation, followed by a dose on Day 5, 14 and 28 after transplant Qty: 4 doses

Documentation Required (Note: Send all labs, must include specific labs listed here)

Labs (TB and Hep B) Insurance Card (front and back) Current Medications History/Progress Notes

Pre-Medication Order (given 30 min before each infusion)

acetaminophen (Tylenol)	500mg	650mg	1000mg PO	diphenhydrAMINE (Benadryl)	25mg	50mg /	PO	IV
cetirizine (ZyrTEC)	10mg PO			methylPREDNISolone (SOLU-Medrol)	40mg IV		125mg IV	
loratadine (Claritin)	10mg PO			hydrocortisone (Solu-CORTEF)	100mg IV			

Other: _____
Dose: _____ Route: _____ Frequency: _____

Line/Port Use Orders

RN to start/discontinue peripheral IV or access/de-access port

RN to administer 3-10 mL normal saline with heparin 100 units/mL 1-5 mL per SASH method

Do NOT administer heparin to this patient

Additional orders: _____

Adverse Reaction and Anaphylaxis Orders

Standard anaphylaxis kit (EPINEPHrine 0.3 mg IM PRN for severe reactions x 1, may repeat once within 5-15 minutes for a maximum of 2 doses; Benadryl 50 mg PO PRN for mild-moderate reactions x1; Benadryl 50 mg IV PRN for mild-moderate reactions x1)

Additional orders: _____

Special Instructions (Prior therapy, treatment dates, and reasons for d/c)**Provider Name**

Order is valid for 1 year from date of signature and refills will be provided to cover the duration of treatment unless otherwise indicated.

o 818.390.9696 aleracare.com

Provider Signature**Date**

Check here if this is a stat order

advanced home infusion™